

Napoli: DSF-T-29/93, INFN-NA-IV-29/93

Syracuse: SU-4240-550

October 1993

FINITE APPROXIMATIONS TO QUANTUM PHYSICS: QUANTUM POINTS AND THEIR BUNDLES

A.P. Balachandran, G. Bimonte, E. Ercolessi

and

P. Teotonio-Sobrinho

*Department of Physics, Syracuse University,
Syracuse, NY 13244-1130*

Abstract

There exists a physically well motivated method for approximating manifolds by certain topological spaces with a finite or a countable set of points. These spaces, which are partially ordered sets (posets) have the power to effectively reproduce important topological features of continuum physics like winding numbers and fractional statistics, and that too often with just a few points. In this work, we develop the essential tools for doing quantum physics on posets. The poset approach to covering space quantization, soliton physics, gauge theories and the Dirac equation are discussed with emphasis on physically important topological aspects. These ideas are illustrated by simple examples like the covering space quantization of a particle on a circle, and the sine-Gordon solitons.

1. Introduction

All known successful fundamental models for the description of nature are based on continuum physics. In the latter, we typically assume that spacetime is a manifold and describe observables by classically smooth structures painted on this background. Such models have given us spectacularly faithful accounts of nature after quantisation, agreement between theory and experiment for electron $g - 2$ being about one part in 10^8 [1]. There is at present no empirical evidence whatsoever that spacetime at a fundamental level is anything but a continuum.

Very good reasons nevertheless exist for studying discrete physics. Natural laws are often exceedingly complex and their quantitative investigation involves their discretisation and adaptation to machines. Also discrete models have many times provided us with soluble systems which illustrate physical principles, or approximate the more accurate continuum theory in a satisfactory manner, well known examples being tight binding [2], Ising [3] and Hubbard models [4] in chemistry and condensed matter theory.

As another reason for the study of discrete physics, mention may also be made of suggestions that spacetime has a discrete structure at the Planck scale [5].

In this paper, we address a serious limitation of conventional discretisations of continuum theories caused by their naive treatment of topology. In lattice QCD [6] for example, spacetime topology is trivial with each point forming an open set and bears no resemblance to the continuum topology. There are similar defects, particularly harmful for topological studies, in discretised investigations of vortices in Higgs models [7] or of solitons in Skyrme physics [8]. No nontrivial winding numbers for instance exist in the usual finite approximations to the Skyrme soliton while it is virtually impossible to display the qualitative bundle theoretical features of monopoles [8] in trivial discretisations.

Some time ago, Sorkin [9] revived and extended an old method [10] for the approxi-

mation of continuous spaces by finite or countable partially ordered sets (posets). This approximation, as we shall see, is physically well motivated. Further the topology of posets, derived from the continuum and approximating the latter, is not simple minded. In particular, the homotopy and homology groups π_n and H_n of posets [9, 10, 11, 12] are generically nontrivial if that is the case for their continuum limits. These approximations are often so good that the continuum π_n and H_n are reproduced with just a few points [13].

In this paper, we initiate the development of tools for doing quantum and soliton physics on posets. An effort has been made to keep the discussion elementary even at the expense of avoiding important proofs or physically significant examples. In Section 2, earlier research and especially Sorkin's work are summarised and illustrated using low dimensional manifolds. Their approximations by posets, and the representation of the latter by Hasse diagrams [9, 12] are also explained. Then to show how accurate the poset approximation can be, we examine π_1 for a four point poset approximation to the circle S^1 and argue that it is $\mathbf{Z}$, the same as for S^1 .

Section 3 introduces a few of the basic tools, and in particular the poset Laplacian Δ , essential for quantum physics. The construction of the latter involves the definition first of a certain boundary operator ∂ on posets [11] and of its dual d , much as in homology and cohomology theories. Intruducing a scalar product on functions of posets, the adjoint $d^\dagger$ or δ of d can then be defined, Δ being just $-d\delta - \delta d$ exactly as for continuum [14]. Energy levels of a 'free' particle on circle posets are also briefly examined as an illustration.

It is well known that the energy eigenstates of a free particle on S^1 can change by a phase when one goes around S^1 , this phase being occasionally explained with the help of a magnetic flux threading S^1 [15]. For manifolds, a general approach to such possibilities involves regarding wave functions as functions on the universal covering space $\overline{Q}$ of the configuration space $Q = \overline{Q}/\pi_1(Q)$ [$\overline{Q}$ being $\mathbf{R}^1$ for $Q = S^1$] which transform by a unitary

irreducible representation (UIR) of $\pi_1(Q)$ [$\pi_1(S^1)$ being $\mathbf{Z}$] [8]. We show in Section 4 that similar ideas work well for posets too, every poset P having another simply connected poset $\overline{P}$ as its universal cover with $P = \overline{P}/\pi_1(P)$. There is also a Laplacian $\overline{\Delta}$ on $\overline{P}$. For circle posets, the spectrum of $\overline{\Delta}$ for wave functions transforming by a UIR of $\pi_1(P)$ is worked out and shown to reproduce the flux effect previously alluded to.

Section 5 deals with topological solitons. The space F of soliton fields consists of smooth maps of a domain space D to a target space T in continuum physics. We discretise D and T to posets $P(D)$ and $P(T)$ and consider instead the space of continuous maps of $P(D)$ to $P(T)$ as approximations of F . We argue that this space $P^{(0)}(F)$ has topological sectors and can describe topological solitons, and can also be converted to a poset. So quantum physics can be done using $P^{(0)}(F)$. The nature of $P^{(0)}(F)$ is illustrated for $D = T = S^1$, the appropriate choice for the sine-Gordon model [8].

This introductory article does not discuss several important topics like Dirac equation and gauge theories on posets, and the computerisation of poset calculations, until Section 6. The latter concludes the paper by outlining our progress in their study.

2. The Finite Approximation

Let Q be a continuous topological space like for example the sphere S^N or the Euclidean space $\mathbf{R}^N$. Experiments are never so accurate that they can detect events associated with points of Q , rather they will only detect events as occurring in certain sets O_λ . It is therefore natural to identify any two points x, y of Q if every set O_λ containing either point contains the other too. Let us assume that the sets O_λ cover Q ,

$$Q = \bigcup_{\lambda} O_{\lambda} \tag{2.1}$$

and write $x \sim y$ if x and y are not separated or distinguished by O_λ in the sense above:

$$x \sim y \text{ means } x \in O_{\lambda} \Leftrightarrow y \in O_{\lambda} . \tag{2.2}$$

Then $\sim$ is an equivalence relation, and it is reasonable to replace Q by $Q/\sim \equiv P(Q)$ to reflect the coarseness of observations. It is this space, obtained by identifying equivalent points, that will be our approximation for Q .

We assume that the number of O_λ is finite when Q is compact so that $P(Q)$ is an approximation to Q by a finite set in this case. When Q is not compact, we assume instead that each point has a neighbourhood intersected by only finitely many O_λ so that $P(Q)$ is a “finitary” approximation to Q [9]. We also assume that each O_λ is open [16] and that

$$\mathcal{U} = \{O_\lambda\} \tag{2.3}$$

is a topology for Q . This implies that $O_\lambda \cup O_\mu$ and $O_\lambda \cap O_\mu \in \mathcal{U}$ if $O_{\lambda,\mu} \in \mathcal{U}$. Now experiments can isolate events in $O_\lambda \cup O_\mu$ and $O_\lambda \cap O_\mu$ if they can do so in O_λ and O_μ separately, the former by detecting an event in either O_λ or O_μ , and the latter by detecting it in both O_λ and O_μ . The hypothesis that $\mathcal{U}$ is a topology is thus conceptually consistent.

These assumptions allow us to isolate events in certain sets of the form $O_\lambda \setminus [O_\lambda \cap O_\mu]$ which may not be open. This means that there are in general points in $P(Q)$ coming from sets which are not open in Q .

In the notation we employ, if $P(Q)$ has N points, we sometimes denote it by $P_N(Q)$.

Let us illustrate these considerations for a cover of $Q = S^1$ by four open sets as in Fig. 1(a). In that figure, $O_{1,3} \subset O_2 \cap O_4$. Figure 1(b) shows the corresponding discrete space $P_4(S^1)$, the points x_i being the images of sets in S^1 . The map $S^1 \rightarrow P_4(S^1)$ is given by

$$\begin{aligned} O_1 &\rightarrow x_1, & O_2 \setminus [O_2 \cap O_4] &\rightarrow x_2, \\ O_3 &\rightarrow x_3, & O_4 \setminus [O_2 \cap O_4] &\rightarrow x_4. \end{aligned} \tag{2.4}$$

Now $P(Q)$ inherits the quotient topology from Q [16]. It is defined as follows. Let Φ be the map from Q to $P(Q)$ obtained by identifying equivalent points. An example of

Φ is given by (2.4). In the quotient topology, a set in $P(Q)$ is declared to be open if its inverse image for Φ is open in Q . It is the finest topology compatible with the continuity of Φ . We adopt it hereafter as the topology for $P(Q)$.

This topology for $P_4(S^1)$ can be read off from Fig.1, the open sets being

$$\{x_1\}, \{x_3\}, \{x_1, x_2, x_3\}, \{x_1, x_4, x_3\}, \quad (2.5)$$

and their unions and intersections (an arbitrary number of the latter being allowed, $P_4(S^1)$ being finite).

A partial order $\preceq$ [10, 11, 12] can be introduced in $P(Q)$ by declaring that $x \preceq y$ if every open set containing y contains also x . For $P_4(S^1)$, this order reads

$$x_1 \preceq x_2, \quad x_1 \preceq x_4; \quad x_3 \preceq x_2, \quad x_3 \preceq x_4, \quad (2.6)$$

where we have omitted writing the relations $x_j \preceq x_j$.

Later, we will write $x \prec y$ to indicate that $x \preceq y$ and $x \neq y$.

It is to be observed that if $x \preceq y$, then the sequence $x, x, x, \dots$ converges to y by the usual definition of convergence. So we may also write $x \rightarrow y$ if $x \preceq y$.

In a Hausdorff space [16], there are open sets O_x and O_y for any two distinct points x and y such that $O_x \cap O_y = \emptyset$. A finite Hausdorff space has discrete topology with each point being an open set. So $P(Q)$ is not Hausdorff. But it is what is called T_0 , where for any two distinct points, there is an open set containing at least one of these points and not the other. For x_1 and x_2 of $P_4(S^1)$, the open set $\{x_1\}$ contains x_1 and not x_2 , but there is no open set containing x_2 and not x_1 .

Any poset can be represented by a Hasse diagram constructed using the following rules: 1) If $x \prec y$, then y is higher than x . 2) If $x \prec y$ and there is no z such that $x \prec z \prec y$, then x and y are connected by a line called a link.

In case 2), y is said to cover x .

The Hasse diagram for $P_4(S^1)$ is shown in Fig. 2.

The smallest open set O_x containing x consists of all y converging to x ($y \rightarrow x$) [so that the closure of the singleton set $\{y\}$ contains x]. In the Hasse diagram, it consists of x and all points we encounter as we travel along links from x to the bottom. In Fig. 2, this rule gives $\{x_1, x_2, x_3\}$ as the smallest open set containing x_2 , just as in (2.5). As another example, consider the Hasse diagram of Fig. 3 for a two-sphere poset $P_6(S^2)$ derived in [9]. Its open sets are generated by

$$\begin{aligned} &\{x_1\}, \{x_3\}, \{x_1, x_2, x_3\}, \{x_1, x_4, x_3\} \\ &\{x_1, x_2, x_5, x_4, x_3\}, \{x_1, x_2, x_6, x_4, x_3\}, \end{aligned} \quad (2.7)$$

by taking unions and intersections.

As one more example, Fig. 4 shows a cover of S^1 by $2N$ open sets O_j and the Hasse diagram of its poset $P_{2N}(S^1)$.

We conclude this Section by arguing that the fundamental group π_1 of $P_4(S^1)$ is $\mathbf{Z}$ [9]. This group is obtained from continuous maps of S^1 to $P_4(S^1)$, or equivalently, of such maps of $[0, 1]$ to $P_4(S^1)$ with the same value at 0 and 1. Figure 5 shows maps like this. The maps shown are continuous, the inverse images of open sets being open [16]. The map in Fig. 5(a) can be deformed to the constant map and has zero winding number. The image in Fig. 5(b) “winds once around” $P_4(S^1)$. The map in this figure has winding number one and is not homotopic to the map in Fig. 5(a). It leads in a conventional way to the generator of $\mathbf{Z}$. $\mathbf{Z}$ is also of course $\pi_1(S^1)$.

3. The Laplacian

We show the construction of the Laplacian Δ for the circle poset $P_{2N}(S^1)$. Its generalization for arbitrary posets will also be indicated.

The numbers ± 1 in Fig. 4(b) are called incidence numbers $I(x_i, x_j)$ [11, 12]. They are attached to each link $x_i - x_j$ where x_i is regarded as less than x_j , $x_i \prec x_j$. We will also

regard $I(x_\alpha, x_\beta)$ as zero if x_β does not cover x_α . The boundary operator ∂ can then be defined by

$$\begin{aligned}\partial x_i &= \sum_{x_j} I(x_i, x_j) x_j \text{ for } i \text{ odd,} \\ \partial x_i &= 0 \text{ for } i \text{ even,}\end{aligned}\tag{3.1}$$

where the sum in (3.1) has a meaning as in homology theory [11]. (3.1) reads, in detail,

$$\partial x_1 = x_2 - x_{2N}, \quad \partial x_3 = x_4 - x_2, \quad \dots \partial x_{2N-1} = x_{2N} - x_{2N-2}, \quad \partial x_{2j} = 0, \quad 1 \leq j \leq N. \tag{3.2}$$

The fundamental property of ∂ ,

$$\partial^2 = 0 \tag{3.3}$$

is trivial in this example.

Next let f_j be the characteristic functions supported at x_j :

$$f_j(x_k) \equiv \langle f_j, x_k \rangle = \delta_{jk}. \tag{3.4}$$

[We will use $\langle f, x \rangle$ and $f(x)$ interchangeably in what follows.] They are a basis for functions on $P_{2N}(S^1)$ with values in $\mathbf{C}$. The boundary operator d on functions is then defined by

$$\langle df_j, x_k \rangle = \langle f_j, \partial x_k \rangle \tag{3.5}$$

or

$$df_2 = f_1 - f_3, \quad df_4 = f_3 - f_5, \quad \dots df_{2N} = f_{2N-1} - f_1, \quad df_{2j-1} = 0, \quad 1 \leq j \leq N. \tag{3.6}$$

It fulfills

$$d^2 = 0. \tag{3.7}$$

(3.6) shows that f_j can be regarded as forming the “dual” poset upside down to Fig.4(b), with f_{2j} at bottom and f_{2j-1} at top, and with the same incidence numbers on links.

The functions on $P_{2N}(S^1)$ turn into a Hilbert space on introducing the scalar product

$$(f_j, f_k) = \delta_{jk}. \quad (3.8)$$

Let $d^\dagger$ be the adjoint of d for (3.8):

$$(d^\dagger f_j, f_k) = (f_j, df_k) \quad \text{or}$$

$$d^\dagger f_1 = f_2 - f_{2N}, \quad d^\dagger f_3 = f_4 - f_2, \dots, d^\dagger f_{2N-1} = f_{2N} - f_{2N-2}, \quad d^\dagger f_{2j} = 0, \quad 1 \leq j \leq N. \quad (3.9)$$

So $d^\dagger$ behaves like ∂ . The Laplacian Δ is

$$\Delta = -dd^\dagger - d^\dagger d \quad (3.10)$$

or

$$\begin{aligned} \Delta f_j &= f_{j+2} + f_{j-2} - 2f_j, \quad \text{for } j \neq 1, 2, (2N-1) \text{ and } 2N, \\ \Delta f_1 &= f_3 + f_{2N-1} - 2f_1, \\ \Delta f_2 &= f_4 + f_{2N} - 2f_2, \\ \Delta f_{2N-1} &= f_1 + f_{2N-3} - 2f_{2N-1}, \\ \Delta f_{2N} &= f_2 + f_{2N-2} - 2f_{2N}. \end{aligned} \quad (3.11)$$

Δ looks like the usual discrete Laplacian for each level, but does not mix the levels.

From analogy to homology theory [11], it is plausible that x_{2j} are to be thought of as corresponding to points of the continuum, and x_{2j-1} as approximations to lines of the continuum connecting x_{2j-2} and x_{2j} . Thus (3.6) can be written as

$$(df)(x_{2j}) = 0, \quad (df)(x_{2j-1}) = f(x_{2j}) - f(x_{2j-2}); \quad (3.12)$$

for any function showing that df at x_{2j-1} corresponds to the integral of df on such a line. Simplicial decompositions of the posets also suggest a similar idea. With these interpretations at hand, one can see that the continuum Laplacian on functions is being

approximated here by the restriction of Δ to functions supported at the top, or at the “rank” zero points, of the poset. [Cf. Section 5.]

The eigenvalues of Δ for $P_{2N}(S^1)$ are

$$\lambda_k = 2(\cos 2k - 1), \quad k = m\frac{\pi}{N}, \quad m = 1, 2, \dots, N. \quad (3.13)$$

For the corresponding eigenfunctions $\varphi^{(k)}$, we have

$$\Delta\varphi^{(k)} = \lambda_k\varphi^{(k)},$$

$$\varphi^{(k)}(x_j) \equiv \varphi_j^{(k)} = e^{ikj}. \quad (3.14)$$

Since $\varphi_{j+2N}^{(k)} = \varphi_j^{(k)}$, they can be regarded as periodic functions on the lattice. They are discrete approximations to plane waves while λ_k are similar approximations to the continuum eigenvalues $-4k^2$. Note that even and odd j separately furnish eigenfunctions for the same eigenvalue.

These considerations can be generalised to any poset by introducing incidence numbers $I(x_\alpha, x_\beta)$ with the following properties [11]:

$$I(x_\alpha, x_\beta) = 0 \quad \text{unless } x_\beta \text{ covers } x_\alpha;$$

$$\sum_{x_\beta} I(x_\alpha, x_\beta) I(x_\beta, x_\gamma) = 0. \quad (3.15)$$

The boundary operator ∂ is then defined by

$$\partial x_\alpha = \sum_{x_\beta} I(x_\alpha, x_\beta) x_\beta,$$

$$\partial x_\alpha = 0 \quad \text{if no point covers } x_\alpha, \quad (3.16)$$

and fulfills

$$\partial^2 = 0. \quad (3.17)$$

The operator d on functions is defined using duality as in (3.5) and fulfills the nilpotency (3.7) because of (3.17). As for the definition of $d^\dagger$ and Δ , we can introduce a scalar

product through

$$(f_j, f_k) = g_{jk} \quad (3.18)$$

where $f_j(x_k) = \delta_{jk}$. $d^\dagger$ is the adjoint of d for this scalar product, while Δ is just (3.10).

It is desirable to restrict g_{jk} somewhat for physical reasons. For example, it is reasonable to suppose that $g_{jk} = 0$ if x_j and x_k approximate surfaces of different dimension in the continuum or are of different rank [cf. Section 5]. Locality is also useful for restricting g_{jk} . We can for example set g_{jk} to zero unless $x_j = x_k$ or there is a point x_l linked to both x_j and x_k . The simplest choice is of course $g_{jk} = \delta_{jk}$.

4. The Universal Cover and the Flux Through the Circle

A charged particle on S^1 with magnetic flux through its middle can be described by the free Hamiltonian, but with “twisted” boundary conditions (BC’s) for energy eigenstates ψ [15]. They (and their derivatives) are not periodic in the coordinate $\varphi \bmod 2\pi$ of S^1 , but change by $e^{i\theta}$ when φ increases by 2π .

It is well known [8] that there is a universal method for the description of such twists. In this general approach, for the preceding example, we regard ψ as functions on the universal covering space $\mathbf{R}^1$ of S^1 . Here if x is the coordinate on $\mathbf{R}^1$, e^{ix} gives the point of S^1 , so that all points $x + 2\pi N$ ($N \in \mathbf{Z}$) of $\mathbf{R}^1$ describe the same point of S^1 . The twist is now got by requiring that $\psi(x + 2\pi) = e^{i\theta}\psi(x)$. This rule is physically sensible, probability densities being single valued on S^1 .

The general method for introducing these twists in continuum physics is as follows [8]. If Q is the configuration space, we let $\overline{Q}$ be its universal covering space. The fundamental group $\pi_1(Q)$ of Q acts freely on $\overline{Q}$ and its quotient $\overline{Q}/\pi_1(Q)$ by this action is Q . Wave functions are obtained from vector valued (one or many component) functions on $\overline{Q}$

transforming by a unitary irreducible representation (UIR) of $\pi_1(Q)$. For $Q = S^1$, $\overline{Q} = \mathbf{R}^1$ and $\pi_1(Q) = \mathbf{Z}$.

We now show that these ideas work for posets as well. Consider first $P_{2N}(S^1)$. Its universal cover is just $P(\mathbf{R}^1)$ shown in Fig. 6. [Its incidence numbers have been obtained from a prescription below.] $\pi_1(P_{2N}(S^1))$ is $\mathbf{Z}$. It acts according to $x_i \rightarrow x_{i+2N}$ and the quotient $P(\mathbf{R}^1)/\mathbf{Z}$ is $P_{2N}(S^1)$.

A systematic approach to find the universal cover $\overline{P}$ of a poset P uses the path space $\mathcal{P}$ as for manifolds [8]. $\mathcal{P}$ consists of continuous maps of $[0, 1]$ to P with the image of 0 being always a fixed point x_0 of P . Let us regard two paths Γ_x and $\Gamma'_x \in \mathcal{P}$ ending at x as equivalent if they are homotopic. A point of $\overline{P}$ is then the equivalence class $[\Gamma_x]$ containing Γ_x . Finally exactly as for manifolds, $\pi_1(P)$ acts by attaching loops to Γ_x at x_0 and $P = \overline{P}/\pi_1(P)$.

Now $\overline{P}$ is also a poset. Its poset structure is got as follows: We assume that $[\Gamma_{x'}]$ covers $[\Gamma_x]$ if a) x' covers x , and b) there is a $\tilde{\Gamma}_{x'}$ in $[\Gamma_{x'}]$ which is a curve $\tilde{\Gamma}_x \in [\Gamma_x]$ followed by the point x' . The incidence number for the link from $[\Gamma_x]$ to $[\Gamma_{x'}]$ is taken to be the same as for the link from x to x' .

Let us finally illustrate how to approximate twisted BC's of S^1 [15] using $P(\mathbf{R}^1)$. The dual d of ∂ can be obtained for Fig. 6 in the usual way. Introducing the scalar product for functions on $P(\mathbf{R}^1)$ through $(f_j, f_k) = \delta_{jk}$ where f_j as usual fulfills $f_j(x_k) = \delta_{jk}$, we can also define $d^\dagger$. The Laplacian Δ for $P(\mathbf{R}^1)$ is then $-dd^\dagger - d^\dagger d$. It is easy to check that

$$\Delta f_j = f_{j+2} + f_{j-2} - 2f_j. \quad (4.19)$$

After imposing the twisted BC's

$$f_{j+2N} = e^{i\theta} f_j, \quad (4.20)$$

the eigenstates $\chi^{(k)}$ supported on the top level and the corresponding eigenvalues λ_k are

found to be given by

$$\begin{aligned}\chi^{(k)}(x_j) &\equiv \chi_j^{(k)} = A^{(k)}e^{ikj} + B^{(k)}e^{-ikj} \\ \lambda_k &= 2(\cos 2k - 1)\end{aligned}\tag{4.21}$$

where $k = m\frac{\pi}{N} + \frac{\theta}{2N}$, $m = 1, 2, 3, \dots, N$. These are obvious approximations to the continuum answers $-4k^2$ for the eigenvalues.

5. Solitons

Topological solitons are characterized by topological invariants such as winding numbers of maps from a manifold D to a manifold T [8]. The existence and properties of these solitons are therefore decisively influenced by the topology of D and T . The use of posets, which can approximate important aspects of topology well, is particularly appropriate for their study.

The ideal method to approximate the space F of smooth maps from D to T would consist in covering each connected component of F by a finite number of open sets. We have not been successful in developing this programme however, F being infinite dimensional. An alternative to this ideal is to substitute D and T by posets $P(D)$ and $P(T)$ and consider continuous maps $P^{(0)}(F)$ from $P(D)$ to $P(T)$, the superscript denoting continuity. We adopt this method here.

An important feature of this approximation can immediately be pointed out. Let us assume that the connected components of F can be labelled by integer valued winding numbers. If M and N are the number of points of $P(D)$ and $P(T)$, then it is impossible to find winding numbers greater than M/N in $P^{(0)}(F)$. This is because it is evidently not possible to cover T by D more often than this ratio. Thus not all continuum solitons can be realized in this approximation with a given M and N .

We now outline our steps for doing quantum physics using $P^{(0)}(F)$. The formalism we develop is an approximation to the field theoretic Schrödinger picture, wave functions being functions of fields. There are two parts to the outline. The first is about the construction of the Laplacian Δ . Δ gives the finite approximation to the continuum kinetic energy. The second part constructs the finite approximation to the gradient energy [which corresponds to an expression like the spatial integral of $(\partial_i \phi \partial^i \phi)$, ϕ being a field.] Of course if there are other potential energy terms in continuum dynamics, they too can be finitely approximated: we see no problem in doing so.

Before going further, we give a convenient characterisation of $P^{(0)}(F)$ which is the set of all continuous functions from $P(D)$ to $P(T)$. By definition, the inverse images of open sets are open for continuous functions. It is easily shown that a function f is continuous iff it preserves order, that is iff $f(x) \preceq f(y)$ whenever $x \preceq y$.

5.1 The Kinetic Energy

The first step in finding Δ is to convert $P^{(0)}(F)$ into a poset. This is readily done: we set $f \preceq g$ if $f(x) \preceq g(x)$ for all $x \in P(D)$.

The next task is to find a ∂ . Now ∂ naturally exists for the space $P(F)$ of all maps from $P(D)$ to $P(T)$ including discontinuous ones. Let us now explain this ∂ , its use for our problem will get clarified later.

A function $f \in P(F)$ is specified by its values $f(i)$ at the points i of $P(D)$ and can thus be written as $(f(1), f(2), \dots, f(M))$. [The points of $P(D)$ can be numbered in any way here.] There being N choices for each entry, $P(F)$ has N^M points.

The points y of $P(T)$ can be assigned a rank $r(y)$ as follows. A point of $P(T)$ is regarded as of rank 0 if it converges to no point, or is a highest point. Let P_1 be the poset got from $P(T)$ by removing all rank zero points and their links. The highest points of P_1 are assigned rank 1. We continue in this way to rank all points.

We will assume that $P(T)$ comes with a boundary operator ∂_T which raises the rank by 1. It is known that such a ∂_T can be defined if $P(T)$ is a “rankable” poset [11]. What this means is the following. Starting from a point y_k of rank k , we can imagine travelling upwards in the Hasse diagram along links till we arrive at a rank zero point. In a rankable poset, the number of links traversed on any such path is always k . Figure 7 shows the Hasse diagram of a poset which is not rankable.

The operator ∂ for $P(F)$ is defined by

$$\begin{aligned} \partial(f(1), f(2), \dots, f(M)) &= (\partial_T f(1), f(2), \dots, f(M)) + (-1)^{r(f(1))} (f(1), \partial_T f(2), \dots, f(M)) + \dots = \\ &= \sum_{j=1}^M (-1)^{\sum_{i < j} r(f(i))} (f(1), f(2), \dots, \partial_T f(j), \dots, f(M)). \end{aligned} \quad (5.1)$$

Clearly, $\partial^2 = 0$ as it should be.

There being topological solitons, $P^{(0)}(F) = \bigcup_k P^{(0)k}(F)$ where $P^{(0)k}(F)$ is a connected topological sector. There is no order relation between elements of $P^{(0)k}(F)$ and $P^{(0)l}(F)$ if $k \neq l$.

Having got ∂ for $P(F)$, we may next try to restrict it to $P^{(0)k}$ for each fixed k . But ∂ on a continuous function will in general involve discontinuous functions, namely those larger than continuous functions. Let $\overline{P}^{(0)k}$ denote the closure of $P^{(0)k}$ obtained by adding all these limit points of $P^{(0)k}$. The ∂ operator of (5.1) is then well defined on $\overline{P}^{(0)k}$.

Figure 8 shows a few elements of $P(F)$ for $D = T = S^1$ and $P(D) = P(T) = P_4(S^1)$. The arrows on the axes indicate the order relations. The x_i on abscissa for example can be identified with those in Fig.2 while y_i are points similar to x_i for the image poset. We remark that since continuous functions preserve order, only those in Figs. 8 (a,b) are continuous.

These figures show a basic danger in working with $\overline{P}^{(0)k}(F)$. The discontinuous function there is contained in both $\overline{P}^{(0)0}$ and $\overline{P}^{(0)1}$ so that different winding number sectors get mixed up with its inclusion. There is thus the risk of losing solitons altogether by

closing $P^{(0)k}$.

We have shown [17] that this difficulty does not arise in this example if $P(T)$ has more than four points. A general result of this kind has also been developed [17].

Having obtained a ∂ , it is now relatively straightforward to find Δ as explained in Section 3.

We further address such issues elsewhere [17, 18, 19].

5.2. The Gradient Energy

The gradient energy in continuum physics involves differentiation with respect to the domain variable whereas ∂ in (5.1) corresponds to differentiation in $P(T)$. We must thus invent a new boundary operator $\hat{\partial}$ to define the gradient energy.

Such a $\hat{\partial}$ is readily available. Let ∂_D be the boundary operator for $P(D)$ (which we assume exists). If $f \in P^{(0)}(F)$ and $x \in D$, let us write $f(x)$ as $\langle f, x \rangle$. $\hat{\partial}$ is then defined by duality:

$$\langle \hat{\partial}f, x \rangle \equiv \langle f, \partial_D x \rangle. \quad (5.2)$$

$\partial_D x$ here is an element of the complex vector space generated by points of D , while f is dual to this space with values in the complex vector space generated by points of T .

Next we introduce a scalar product $(\cdot, \cdot)$ corresponding to T as in (3.18). $\langle \hat{\partial}f, x \rangle$ being a combination of points of T , $(\langle \hat{\partial}f, x \rangle, \langle \hat{\partial}f, x \rangle)$ is well defined and nonnegative and can be identified as being proportional to the gradient energy density. The total gradient energy is obtained by summing over x .

All the tools for studying finite approximations to solitons are now at hand.

6. Concluding Remarks

We have not yet mentioned several important topics for reasons of simplicity. We will conclude the paper by pointing out a few of these topics and our own progress in their study.

1) The first topic concerns the utility of the poset approximation for practical calculations. Definite progress has already been made in its study [19]. Algorithms for generating soliton posets have been devised and tested. Numerical work on sine-Gordon solitons are now under way.

2) Another basic subject for research is the poset variant of the Dirac equation. If d and δ have the conventional meanings of Section 3, it is known that the familiar Dirac equation in Minkowsky space can be obtained from $[d + \delta + m]\psi = 0$ [20]. It is also known that a similar equation on ordinary lattices gives the lattice variant of this equation [20]. As both d and δ make sense on a typical poset, the similar poset equation seems a good starting point to formulate the poset Dirac equation.

3) Progress has been made in the study of gauge theories too. When space- time, and not space alone, is made a poset, we have discovered a natural way to get the Wilson action [6]. We now outline the idea, leaving details to another paper. In the absence of gauge fields, the operator d acts on a function f of $P(Q)$ by a rule like $df(i) = f(j) - f(k)$, with j and k covering i . Its gauge theory version $\mathcal{D}$ can be obtained by first introducing the link variables with values U_{ij} , U_{ik} in the gauged group G , regarded here as unitary matrices acting on a vector space V . With the function f also thought of as valued in V , $\mathcal{D}$ can be defined using the equation $(\mathcal{D}f)(i) = U_{ij}f_j - U_{ik}f_k$. It is a covariant equation as in lattice gauge theories. Assuming that $U_{ji} = U_{ij}^{-1}$, we can also show that $(\mathcal{D}^2f)(i)$ is of the form $[U_{ij}U_{jk}U_{kl}U_{li}]f(i)$. The Wilson action is then just a sum over terms of the form constant $\times \text{Tr}[U_{ij}U_{jk}U_{kl}U_{li}]$.

The poset Hamiltonian formulation for $G = U(1)$ also appears to present no difficulty and can be achieved as follows. $U(1)$ is first approximated by $P_{2N}(U(1)) = P_{2N}(S^1)$. It

is also convenient to build up wave functions from functions on Wilson integrals. For this purpose, we assign $U_{ij} \in P_{2N}(S^1)$ to each link ij of $P(Q)$, the poset for the spatial slice Q , with $U_{ji} = U_{ij}^{-1}$. [Note that points of $P_{2N}(S^1)$ can be given a group structure by associating them with $e^{i\frac{\pi}{N}n}$, $n = 1, 2, \dots, 2N$.] Now consider continuous maps of $[0, 1]$ to $P(Q)$ with 0 always having a fixed image x_0 . By multiplying U_{ij} along the image of $[0, 1]$, we get a map W of $[0, 1]$ to $P_{2N}(S^1)$. It is necessary in our approach that W fullfills suitable continuity conditions which we plan to explain later [17], the choice of U_{ij} being accordingly restricted. The space $P^{(0)}(W)$ of these maps is what replaces the space $P^{(0)}(F)$ of Section 5 in gauge theories. As it can usually be made a poset just as $P^{(0)}(F)$, we can also try building Laplacians and other structures appropriate for it.

There is a natural notion of gauge invariance too here. This is because the group $\mathbf{Z}_N$ acts continuously on $P_{2N}(S^1)$, the response of $x_i (\equiv x_{i+2N})$ in Fig.4(b) to its generator being $x_i \rightarrow x_{i+2}$, and we can think of this group as approximating $U(1)$. Gauge transformations on W can be defined using it as in lattice gauge theories[6].

While some of these ideas work for nonabelian G , we find difficulties in conveniently formulating the notion of gauge invariance for such G . The problem is with the symmetries of the approximating posets $P_k(G)$ of G . It is seldom the case that $P_k(G)$ admits a homeomorphism group G_k with G_k approaching G for large k in some sense. Indeed it is already impossible to find discrete subgroups $SU(2)_k$ of $SU(2)$ which converge to $SU(2)$ for large k . The treatment of nonabelian gauge theories in the poset approach thus requires more work.

Acknowledgements

We are very grateful to L. Chandar for numerous discussions and good suggestions throughout the course of this work. We also thank Ajit Mohan Srivastava for help with the references. This work was supported by the Department of Energy under contract number DE-FG02-ER40231. In addition, G.B. was supported by INFN(Italy) and P.T.S

was partially supported by Capes (Brazil).

References

- [1] R.S. Van Dyck Jr., P.B. Schwinberg and H.G. Dehmelt, Phys. Rev. Letters **59** (1987) 26.
- [2] Cf. J. E. Avron, “Adiabatic Quantum Transport”, in “Anomalies, Phases, Defects, ...”, Edited by M. Bregola, G. Marmo and G. Morandi [Bibliopolis, 1990].
- [3] B.M. McCoy, “The Two-Dimensional Ising Model” [Harvard University Press, 1973]; G. Parisi, “Statistical Field Theory” [Addison-Wesley, 1988].
- [4] For a recent review, see A.P. Balachandran, E. Ercolessi, G. Morandi and A.M. Srivastava, “Hubbard Model and Anyon Superconductivity”, Lectures Notes in Physics, Volume 38 [World Scientific, 1990].
- [5] G. 't Hooft, in “Recent Developments in Gravitation”, M. Levy and S. Deser, eds. [Plenum, 1979] ; L. Bombelli, J. Lee, D. Mayer and R.D. Sorkin, Phys. Rev. Letters **59** (1987) 521; R.D. Sorkin, in *Proceedings of the Third Canadian Conference on General Relativity and Relativistic Astrophysics*, A. Coley, F. Cooperstock and B. Tupper, eds. [World Scientific, 1990] and references therein.
- [6] J.M. Drouffe and C. Itzykson, Phys. Rep. **38** (1978) 134; M. Creutz, “Quarks, Gluons and Lattices” [Cambridge University Press, 1983]; J.B. Kogut, Rev. Mod. Phys. **55** (1983) 785.
- [7] “Superconductivity”, vol. 2, R.D. Parks, ed. [Marcel Dekker, 1969]; A. Jaffe and C. Taubes, “Vortices and Monopoles” [Birkhäuser, 1980]; P.G. de Gennes, “Superconductivity of Metals and Alloys” [Addison-Wesley, 1989]; H. Kleinert, “Gauge Fields in Condensed Matter” [World Scientific, 1989].

- [8] Cf. A.P. Balachandran, G. Marmo, B.S. Skagerstam and A. Stern, “Classical Topology and Quantum States” [World Scientific, 1991].
- [9] R.D. Sorkin, Int. J. Theor. Phys. **30** (1991)923.
- [10] P.S. Aleksandrov, *Combinatorial Topology*, Vol. 1-3 [Greylock, 1960].
- [11] P.J. Hilton and S. Wylie, *Homology Theory* [Cambridge, 1966].
- [12] R.P. Stanley, *Enumerative Combinatorics*, vol.I [Wordsworth and Brooks/ Cole Advanced Books and Software, 1986].
- [13] For another approach to discretization which pays attention to topological features, see A.V. Ivashchenko, Syracuse Preprint SU-GP-93/1-1 (1993).
- [14] J.R. Munkres, *Elementary Differential Topology* [Princeton University Press, 1966].
- [15] Cf. C. Aneziris, A.P. Balachandran, M. Bourdeau, S. Jo, T.R. Ramadas and R.D. Sorkin, Int. J. Mod. Phys. **A4** (1989) 5459.
- [16] Encyclopedic Dictionary of Mathematics, edited by Shôkidu Iyanaga and Yukiyo Kawada, translation reviewed by K.O. May [The M.I.T. Press, 1977], Section 408.
- [17] A.P. Balachandran, G. Bimonte, E. Ercolessi and P. Teotonio-Sobrinho (in preparation).
- [18] For a detailed discussion of Δ for a poset approximation of an interval including the renormalization group properties of its continuum limit, see G. Bimonte, E. Ercolessi and P. Teotonio-Sobrinho, Syracuse University and Napoli preprint SU-4240-548 and DSF-T-28/93,INFN-NA-IV-28/93 (1993).
- [19] A.P. Balachandran, G. Bimonte, E. Ercolessi, F. Lizzi and P. Teotonio-Sobrinho (work in progress)

[20] Cf. J.M. Rabin, Nucl. Phys. **B201**, 315 (1982) and references therein.




Fig.1. (a) shows an open cover for the circle S^1 and (b) the resultant discrete space $P_4(S^1)$. Φ is the map (2.4).




Fig.2. The Hasse diagram for the circle poset $P_4(S^1)$.




Fig.3. The Hasse diagram for the two-sphere poset $P_6(S^2)$.





Fig.4. In (a) is shown a covering of S^1 by open sets O_j with $O_3 = O_2 \cap O_4$, $O_5 = O_4 \cap O_6$, ..., $O_1 = O_{2N} \cap O_2$. (b) is the Hasse diagram of its poset with a choice of incidence numbers on the links.




Fig.5. The figures show continuous maps of S^1 to $P_4(S^1)$. The dark lines correspond to closed intervals of S^1 and the light lines to open intervals. The map in (a) has zero winding number, and the map in (b) has winding number 1.




Fig.6. The universal cover $P(\mathbf{R}^1)$ of $P_{2N}(S^1)$.




Fig.7. The Hasse diagram of a poset which is not rankable. There is only one link between y_1 of rank 1 and y_3 of rank 3. We can arrive at y_0 of rank 0 from y_3 encountering only the two links $y_3 - y_1$ and $y_1 - y_0$.




Fig.8. The figure shows some maps of $P_4(S^1)$ to $P_4(S^1)$. (a) shows the unique continuous winding number 1 map, (b) shows three continuous winding number zero maps I, II, and III and (c) represents a discontinuous function lying in both $\overline{P}^{(0)0}(F)$ and $\overline{P}^{(0)1}(F)$. Note that the maps are given by just the dots, the lines connecting them having no special meaning. Note further that $x_3 \rightarrow x_2$, $y_3 \rightarrow y_2$ although these are not shown in the figures.